

Does electron capture dissociation (ECD) provide quantitative information on the chemical modification of lysine side chains in proteins? The glycation of ubiquitin.

Autorzy

Piotr Stefanowicz

Monika Kijewska

Zbigniew Szewczuk

Rok wydania

2014

Czasopismo

Analytical Chemistry

Numer woluminu

86

Strony

7247-7251

DOI

10.1021/ac501329g

Kolekcja

Naukowa

Język

Angielski

Typ publikacji

Artykuł

Streszczenie

Electron capture dissociation (ECD) as a method of quantitative and qualitative study of glycated ubiquitin was investigated. ECD has been successfully applied for sequencing of modified peptides and assigning glycated Lys residues. By using a hybrid Fourier transform mass spectrometry (FT-MS) system equipped for ECD, a series of multiply glycated ubiquitin ions was observed. Ions of the glycated ubiquitin with a defined number of glucose moieties attached to the protein were isolated by quadrupole and fragmented in the ICR cell by the ECD method. The fragmentation spectrum was dominated by c_n and $(z+1)_n$ ions. The ECD technique was tested for the quantitative analysis of the modified ubiquitin and isomeric glycated peptides (fragments of bovine serum albumin (BSA)). Obtained results indicate that the ECD fragmentation cannot be applied for the quantitative determination of the relative reactivities of respective Lys residues in the ubiquitin.

Adres publiczny

<http://dx.doi.org/10.1021/ac501329g>

Strona internetowa wydawcy

<https://www.acs.org/content/acs/en.html>