Thiophosphorylation of free amino acids and enzyme protein by thiophosphoramidate ions.

Autorzy

Tomasz Ruman

K. Długopolska

A. Jurkiewicz

D. Rut

Tomasz Frączyk

J. Cieśla

A. Leś

Zbigniew Szewczuk

Wojciech Rode

Rok wydania

2010

Czasopismo

Bioorganic Chemistry

Numer woluminu

38

Strony

74-80

DOI

10.1016/j.bioorg.2009.11.002

Kolekcja

Naukowa

Język

Angielski

Streszczenie

In search of an activity-preserving protein thiophosphorylation method, with thymidylate synthase recombinant protein used as a substrate, potassium thiophosphoramidate and diammonium thiophosphoramidate salts in Tris- and ammonium carbonate based buffer solutions were employed, proving to serve as a non-destructive environment. Using potassium phosphoramidate or diammonium thiophosphoramidate, a series of phosphorylated and thiophosphorylated amino acid derivatives was prepared, helping, together with computational (using density functional theory, DFT) estimation of ^{31}P NMR chemical shifts, to assign thiophosphorylated protein NMR resonances and prove the presence of thiophosphorylated lysine, serine and histidine moieties. Methods useful for prediction of ^{31}P NMR chemical shifts of thiophosphorylated amino acid moieties, and thiophosphates in general, are also presented. The preliminary results obtained from trypsin digestion of enzyme shows peak at *m/z* 1825.805 which is in perfect agreement with the simulated isotopic pattern distributions for monothiophosphate of TVQQQVHLNQDEYK where thiophosphate moiety is attached to histidine (His^{26}) or lysine (Lys^{33}) side-chain.

Słowa kluczowe

NMR, Thiophosphate, Tiophosphoramidate, Thiophosphorylation, Thymidylate synthase

Adres publiczny

https://doi.org/10.1016/j.bioorg.2009.11.002

Strona internetowa wydawcy

http://www.elsevier.com

Typ publikacji

Artykuł