

## Supramolecular stores of acetic acid—an alternative for metal-organic frameworks.

### Autorzy

Danuta Dobrzyńska  
M. Litwin  
Lucjan B. Jerzykiewicz  
A. Matraszek

### Rok wydania

2009

### Czasopismo

Journal of Hazardous  
Materials

### Numer woluminu

169

### Strony

1040-1044

### DOI

10.1016/j.jhazmat.2009.04.041

### Kolekcja

Naukowa

### Język

Angielski

### Typ publikacji

Artykuł

### Streszczenie

A series of new complexes:  $[M(\text{quin-2-c})_2(\text{H}_2\text{O})_2] \cdot 4\text{CH}_3\text{COOH}$  ( $M = \text{Mn}^{2+}$ ,  $\text{Co}^{2+}$  or  $\text{Ni}^{2+}$ , quin-2-c is quinoline-2-carboxylate ion) have been synthesized and characterized by X-ray single crystal study. The crystals of the complexes reveal very interesting 1D structures comprising acetic acid molecules. The processes of re- and desolvation of acetic acid by the manganese complex in mild conditions have been studied. The de- and resolvated materials were characterized by elemental analysis, IR spectroscopy and XRD study. The results show that binding of acetic acid is reversible and stoichiometric.

### Słowa kluczowe

Acetic acid, Supramolecular network, X-ray study, Quinoline-2-carboxylate ligand

### Adres publiczny

<https://doi.org/10.1016/j.jhazmat.2009.04.041>

### Strona internetowa wydawcy

<http://www.elsevier.com>