Uniwersytet Wrocławski

Repozytorium

Plik pochodzi z repozytorium Wydziału Chemii Uniwersytetu Wrocławskiego

Building 1D lanthanide chains and non-symmetrical [Ln_2] "triple-decker" clusters using salen-type ligands: magnetic cooling and relaxation phenomena.

Autorzy

Angelos B. Canaj

Miłosz Siczek

Marta Otręba

Tadeusz Lis

Giulia Lorusso

Marco Evangelisti

Constantinos J. Milios

Rok wydania

2016

Czasopismo

Dalton Transactions

Numer woluminu

45

Strony

18591-18602

DOI

10.1039/c6dt02932a

Kolekcja

Naukowa

Język

Angielski

Typ publikacji

Artykuł

Streszczenie

A solvothermal reaction between $Ln(NO_3)_3{\cdot}6H_2O$ (Ln: Gd, Tb and Dy), 2-hydroxy-1-naphthaldehyde, 2-OH-naphth, and ethylenediamine, en, in MeOH in the presence of a base, NEt_3, led to the formation of the 1D coordination polymers $[Ln(L)(MeO)(MeOH)_{0.5}]_n{\cdot}MeOH$ (Ln = Gd (**1**·MeOH), Tb(**2**), Dy (**3**·MeOH); H_2L = 1,1′-((1*E*,1′*E*)-(ethane-1,2-diylbis(azanylylidene))bis(methanylylidene))bis(naphthalen-2-ol), the Schiff-base ligand derived from the condensation of 2-OH-naphth and en), while a similar reaction in an excess of NaN_3 yielded 1D coordination polymers $[Ln(L)(N_3)_{0.75}(MeO)_{0.25}(MeOH)]_n$ (Ln = Gd (**4**), Tb (**5**), Dy (**6**)). Finally, upon replacing ethylenediamine with *o*-phenylenediamine, *o*-phen, we managed to isolate the discrete dimers $[Dy_2(L')_3(MeOH)]{\cdot}2MeOH$ (**7**·2MeOH) and $[Gd_2(L')_3(MeOH)]{\cdot}2MeOH$ (**8**·2MeOH) (H_2L' = 1,1′-((1*E*,1′*E*)-(1,2-phenylenebis(azanylylidene))bis(methanylylidene))bis(naphthalen-2-ol), the Schiff-base ligand from the condensation of 2-OH-naphth and *o*-phen). Polymers **1–3** describe one-dimensional chains, containing alternating seven- and eight-coordinate Ln^{III} metal centers, polymers **4–6** contain eight-coordinate lanthanide ions, while in both **7** and **8** the two Ln^{III} centers are eight- and seven-coordinate, adopting square antiprismatic and “piano-stool” geometry, respectively. The magnetocaloric properties of the three Gd^{III} analogues were determined from magnetic measurements, yielding the magnetic entropy change $-\Delta S_m$ = 21.8, 23.0 and 16.0 J kg^{-1} K^{-1} at T = 3.0 K on demagnetization of 7 T to 0, for **1**, **4** and **8**, respectively. The study of the magnetic properties also revealed that all three Dy^{III} analogues (**3**, **6** and **7**) display out-of-phase signals, therefore suggesting slow magnetic relaxation, while such behaviour was not established in the Tb^{III} analogues.

Adres publiczny

http://dx.doi.org/10.1039/c6dt02932a

Strona internetowa wydawcy

https://www.rsc.org/

Plik został wygenerowany dnia 2026-11-13 02:37:39

Adres w repozytorium https://old.chem.uni.wroc.pl/pl/repozytorium/ZwwJnNG.