

Structure-based design, synthesis, and evaluation of the biological activity of novel phosphoroorganic small molecule IAP antagonists.

Autorzy

Agnieszka Łupicka-Słowik

Mateusz Psurski

Renata Grzywa

Monika Cuprych

Jarosław Ciekot

Waldemar Goldeman

Elżbieta Wojaczyńska

Jacek Wojaczyński

Józef Oleksyszyn

Marcin Sieńczyk

Rok wydania

2020

Czasopismo

Investigational New Drugs

Numer woluminu

38

Strony

1350-1364

DOI

10.1007/s10637-020-00923-4

Kolekcja

Naukowa

Język

Angielski

Streszczenie

One of the strategies employed by novel anticancer therapies is to put the process of apoptosis back on track by blocking the interaction between inhibitor of apoptosis proteins (IAPs) and caspases. The activity of caspases is modulated by the caspases themselves in a caspase/procaspase proteolytic cascade and by their interaction with IAPs. Caspases can be released from the inhibitory influence of IAPs by proapoptotic proteins such as secondary mitochondrial activator of caspases (Smac) that share an IAP binding motif (IBM). The main purpose of the present study was the design and synthesis of phosphorus-based peptidyl antagonists of IAPs that mimic the endogenous Smac protein, which blocks the interaction between IAPs and caspases. Based on the structure of the IAP antagonist and recently reported thiadiazole derivatives, we designed and evaluated the biochemical properties of a series of phosphonic peptides bearing the N-Me-Ala-Val/Chg-Pro-OH motif (Chg: cyclohexylglycine). The ability of the obtained compounds to interact with the binding groove of the X-linked inhibitor of apoptosis protein baculovirus inhibitor of apoptosis protein repeat (XIAP BIR3) domain was examined by a fluorescence polarization assay, while their potential to induce autoubiquitination followed by proteasomal degradation of cellular IAP1 was examined using the MDA-MB-231 breast cancer cell line. The highest potency against BIR3 was observed among peptides containing C-terminal phosphonic phenylalanine analogs, which displayed nanomolar K_i values. Their antiproliferative potential as well as their proapoptotic action, manifested by an increase in caspase-3 activity, was examined using various cell lines.

Słowa kluczowe

IAP antagonist, smac mimetic, apoptosis, phosphoroorganic

Typ publikacji

Artykuł

Licencja otwartego dostępu

CC-BY

Licencja na prawach której można swobodnie kopiować, rozprowadzać, zmieniać i remiksować objęty prawem autorskim utwór (Utwór-przedmiot prawa autorskiego) pod warunkiem podania imienia i nazwiska autora utworu pierwotnego oraz źródła pochodzenia utworu.

Pełny tekst licencji:

<https://creativecommons.org/licenses/by/3.0/pl/legalcode>

Adres publiczny

<http://dx.doi.org/10.1007/s10637-020-00923-4>

Strona internetowa wydawcy

<http://link.springer.com>

Plik został wygenerowany dnia 2026-10-03 09:18:57

Adres w repozytorium https://old.chem.uni.wroc.pl/pl/repozytorium/c_JD8W9.