

A hexapeptide VTKFYF from C-terminal part of interleukin-1 receptor antagonist, an inhibitor of IL- 1-IL-1 receptor interaction.

Autorzy

Zbigniew Wieczorek

Alicja Kluczyk

Jacek J. Słoń

Ignacy Z. Siemion

Rok wydania

1997

Czasopismo

Polish Journal of
Pharmacology

Numer woluminu

49

Strony

107-117

Kolekcja

Naukowa

Język

Angielski

Typ publikacji

Artykuł

Streszczenie

In order to find the low-molecular-weight interleukin-1 (IL-1) inhibitors we synthesized a series of peptides, derived from three regions of interleukin-1 receptor antagonist (IL-1ra): N-terminal (residues 5-9), central (90-98) and C-terminal (143-148). The decision was based on the thorough analysis of structural and functional properties of IL-1 proteins and the resemblance of some fragments of IL-1ra to well-known immunomodulators, like thymopentin and tuftsin. The competition between our peptides and IL-1 was measured as the inhibition of IL-1 induced IL-2 production in LBRM/CTLL cell line system. The activity of tuftsin (TKPR), a peptide immunomodulator derived from IgG molecule, was also examined. All peptides presented some activity, although the most interesting results (when the range of activity and dose-dependence were taken into account) were obtained for tuftsin and peptide VTKFYF from the C-terminal part of IL-1ra, which is in agreement with the latest reports on the structure of IL-1ra-receptor complex.

Strona internetowa wydawcy

<http://if-pan.krakow.pl/en/>