

Chiral macrocyclic lanthanide complexes derived from (1*R*,2*R*)-1,2-diphenylethylenediamine and 2,6-diformylpyridine.

Autorzy

Jarosław Mazurek

Jerzy Lisowski

Rok wydania

2003

Czasopismo

Polyhedron

Numer woluminu

22

Strony

2877-2883

DOI

10.1016/S0277-
5387(03)00406-6

Kolekcja

Naukowa

Język

Angielski

Typ publikacji

Artykuł

Streszczenie

The enantiopure complexes $[\text{LnL}]\text{Cl}_3 \cdot n\text{H}_2\text{O}$ ($\text{Ln} = \text{La}^{3+}$, Ce^{3+} and Eu^{3+}) of the new chiral macrocycle derived from (1*R*,2*R*)-1,2-diphenylethylenediamine and 2,6-diformylpyridine have been synthesised. The complexes have been characterised by NMR spectroscopy and mass spectrometry. The ^1H and ^{13}C signal assignment has been based on COSY, NOESY and HMQC measurements. The X-ray crystal structure of the $[\text{CeL}]\text{Cl}_3 \cdot 4\text{H}_2\text{O}$ complex has been determined. The cerium(III) ion is coordinated by six nitrogen atoms of the macrocyclic ligand and three chloride anions. The macrocycle in this complex adopts a relatively flat, twist-bent conformation. The enantiopure La^{3+} , Ce^{3+} and Eu^{3+} complexes of the new chiral macrocycle derived from (1*R*,2*R*)-1,2-diphenylethylenediamine and 2,6-diformylpyridine have been obtained in template condensation. The complexes have been characterised by 2D NMR spectroscopy and mass spectrometry. The X-ray crystal structure of the cerium(III) complex shows the Ce(III) ion coordinated by six nitrogen atoms of the macrocyclic ligand and three chloride anions. The macrocycle in this complex adopts a relatively flat, twist-bent conformation.

Słowa kluczowe

Macrocycles, Lanthanide complexes, Chiral complexes, NMR

Adres publiczny

[https://doi.org/10.1016/S0277-5387\(03\)00406-6](https://doi.org/10.1016/S0277-5387(03)00406-6)

Strona internetowa wydawcy

<http://www.elsevier.com>